

Due applicazioni della fotometria solare allo studio della qualità dell'aria: determinazione del diossido di azoto e delle proprietà ottiche degli aerosol

H. Diémoz ^{1,2} A. M. Siani ² M. Campanelli ³

¹ARPA Valle d'Aosta

²Sapienza Università di Roma

³ISAC-CNR

Convegno nazionale AgF - 8 giugno 2012

1 Introduction

2 Nitrogen dioxide

3 Aerosol properties


4 Conclusions

Introduction

Bouguer-Lambert-Beer law

$$dI = -K_{\text{ext}} I \rho ds \quad (1)$$

$$[K_{\text{ext}}] = \frac{m^2}{kg} \quad (2)$$



[Wikipedia]

Introduction

NO₂ in the atmosphere

Troposphere

- mostly produced from high temperature processes
- one of the most important precursors of tropospheric ozone
- controls the OH radical
- criteria pollutant (very reactive, dangerous for human health: highly oxidant and carcinogenic)
- related to acid rain (HNO₃ is very soluble in water)
- locally contributes to radiative forcing

Stratosphere

- catalyst of ozone destruction (25÷40 km, Crutzen)
 - ▶ ~50% of ozone removal is due to NO₂
- moderates the ozone loss (10÷25 km), converting active chlorine and hydrogen into reservoir species

Introduction

NO₂ in the atmosphere

Troposphere


- mostly produced from high temperature processes
- one of the most important precursors of tropospheric ozone
- controls the OH radical
- criteria pollutant (very reactive, dangerous for human health: highly oxidant and carcinogenic)
- related to acid rain (HNO₃ is very soluble in water)
- locally contributes to radiative forcing

Stratosphere

- catalyst of ozone destruction (25÷40 km, Crutzen)
 - ▶ ~50% of ozone removal is due to NO₂
- moderates the ozone loss (10÷25 km), converting active chlorine and hydrogen into reservoir species


Introduction

NO₂ in the atmosphere



Introduction


NO₂ in the atmosphere



[SCIAMACHY]

Introduction

Aerosols




THE POWER OF POLLUTION

Aerosols — tiny particles from pollution, volcanoes, dust and other sources — can reflect or absorb sunlight directly, or seed cloud droplets and brighten clouds. New climate models suggest that aerosols and clouds can have bigger than expected influences.

[Nature]

Introduction

Aerosols



[Nature]

Introduction

Solar photometry represents a valuable, alternative approach to air quality monitoring!



[ARPA VdA]

1 Introduction

2 Nitrogen dioxide

3 Aerosol properties

4 Conclusions

Aim of present research

- employ a Brewer MKIV spectrophotometer to accurately measure total NO_2 content (and other species as well?) in the visible range
 - ▶ accurately characterize Brewer #066
 - ▶ improve/redesign the retrieval algorithm (new cross-sections, better techniques, ...)
 - ▶ calibrate Brewer #066 ←no calibrated standard MKIV for NO_2 measurements is available worldwide
 - ▶ compare the results of ZS and DS techniques using MKIV
 - ▶ satellite and ground-based instruments intercomparisons?


Nitrogen dioxide

Aim of present research

- employ a Brewer MKIV spectrophotometer to accurately measure total NO₂ content (and other species as well?) in the visible range
 - ▶ accurately characterize Brewer #066
 - ▶ improve/redesign the retrieval algorithm (new cross-sections, better techniques, ...)
 - ▶ calibrate Brewer #066 ←no calibrated standard MKIV for NO₂ measurements is available worldwide
 - ▶ compare the results of ZS and DS techniques using MKIV
 - ▶ satellite and ground-based instruments intercomparisons?


Nitrogen dioxide

Preliminary analysis on solar spectra (Varimax PCA)



Nitrogen dioxide


The Brewer spectrophotometer



Mostly used for ozone measurements (UV)

Nitrogen dioxide


The Brewer spectrophotometer



[Kipp&Zonen]

Nitrogen dioxide

The Brewer spectrophotometer



[Kipp&Zonen]

Nitrogen dioxide

The Brewer spectrophotometer


Characterization

- Focus in the visible range
- Wavelength-steps dispersion function
- Filter transmission in the visible
- Internal polarization
- Straylight
- Temperature dependence

New routines were developed

Nitrogen dioxide

Calibration of the Brewer spectrophotometer



Brewer #066 at Plateau Rosa (3500 m asl), July 2011

Nitrogen dioxide

The classical Brewer algorithm

$$\begin{aligned} F_{corr}(\lambda_i) = & ETC(\lambda_i) - [K_{NO_2}(\lambda_i)\mu_{NO_2}X_{NO_2} \\ & + K_{O_3}(\lambda_i)\mu_{O_3}X_{O_3} \\ & + \delta_{aer}(\lambda_i)\mu_{aer}] \end{aligned} \quad (3)$$

- the contribution of other atmospheric species is assumed to be negligible
- F_{corr} takes into account the attenuation of the filters and Rayleigh scattering

Nitrogen dioxide

The classical Brewer algorithm

$$\Sigma \gamma_i F_i = \Sigma \gamma_i ETC_i - \mu_{NO_2} X_{NO_2} (\Sigma \gamma_i K_i) \quad (4)$$

$$MS_9 = NB_1 + M_2 MS_{11} NA_1 \quad (5)$$

- only NO_2 contribution remains if weighting coefficients are correctly calculated
- the retrieved NO_2 content depends on the NO_2 X-secs
- weighting coefficients depend on the X-secs used for species different from NO_2
- effective X-secs strongly depends on operational wavelengths and shape of the transmission function

Nitrogen dioxide

The classical Brewer algorithm

$$\begin{pmatrix} \sigma_{O_3}(\lambda_1) & \sigma_{O_3}(\lambda_2) & \sigma_{O_3}(\lambda_3) & \sigma_{O_3}(\lambda_4) & \sigma_{O_3}(\lambda_5) \\ \beta_{Ray}(\lambda_1) & \beta_{Ray}(\lambda_2) & \beta_{Ray}(\lambda_3) & \beta_{Ray}(\lambda_4) & \beta_{Ray}(\lambda_5) \\ \delta_{aer}(\lambda_1) & \delta_{aer}(\lambda_2) & \delta_{aer}(\lambda_3) & \delta_{aer}(\lambda_4) & \delta_{aer}(\lambda_5) \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \gamma_5 \end{pmatrix} = \mathbf{0}_{K_{4,1}}$$

i.e.

$$\begin{cases} \vec{\gamma} \perp \vec{\sigma}_{O_3} \\ \vec{\gamma} \perp \vec{\beta}_{Ray} \\ \vec{\gamma} \perp \vec{\delta}_{aer} \\ \vec{\gamma} \cdot (1, 1, 1, 1, 1) = 0 \end{cases}$$

i.e.

$\vec{\gamma} \in \ker(\mathbf{A}) \rightarrow$ only 1 dimension left (normalization)

Nitrogen dioxide

The classical Brewer algorithm

$$\begin{pmatrix} \sigma_{O_3}(\lambda_1) & \sigma_{O_3}(\lambda_2) & \sigma_{O_3}(\lambda_3) & \sigma_{O_3}(\lambda_4) & \sigma_{O_3}(\lambda_5) \\ \beta_{Ray}(\lambda_1) & \beta_{Ray}(\lambda_2) & \beta_{Ray}(\lambda_3) & \beta_{Ray}(\lambda_4) & \beta_{Ray}(\lambda_5) \\ \delta_{aer}(\lambda_1) & \delta_{aer}(\lambda_2) & \delta_{aer}(\lambda_3) & \delta_{aer}(\lambda_4) & \delta_{aer}(\lambda_5) \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \gamma_5 \end{pmatrix} = \mathbf{0}_{K_{4,1}}$$

i.e.

$$\begin{cases} \vec{\gamma} \perp \vec{\sigma_{O_3}} \\ \vec{\gamma} \perp \vec{\beta_{Ray}} \\ \vec{\gamma} \perp \vec{\delta_{aer}} \\ \vec{\gamma} \cdot (1, 1, 1, 1, 1) = 0 \end{cases}$$

i.e.

$\vec{\gamma} \in \ker(\mathbf{A}) \rightarrow$ only 1 dimension left (normalization)

Nitrogen dioxide

The classical Brewer algorithm

$$\begin{pmatrix} \sigma_{O_3}(\lambda_1) & \sigma_{O_3}(\lambda_2) & \sigma_{O_3}(\lambda_3) & \sigma_{O_3}(\lambda_4) & \sigma_{O_3}(\lambda_5) \\ \beta_{Ray}(\lambda_1) & \beta_{Ray}(\lambda_2) & \beta_{Ray}(\lambda_3) & \beta_{Ray}(\lambda_4) & \beta_{Ray}(\lambda_5) \\ \delta_{aer}(\lambda_1) & \delta_{aer}(\lambda_2) & \delta_{aer}(\lambda_3) & \delta_{aer}(\lambda_4) & \delta_{aer}(\lambda_5) \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \gamma_5 \end{pmatrix} = \mathbf{0}_{K_{4,1}}$$

i.e.

$$\begin{cases} \vec{\gamma} \perp \vec{\sigma_{O_3}} \\ \vec{\gamma} \perp \vec{\beta_{Ray}} \\ \vec{\gamma} \perp \vec{\delta_{aer}} \\ \vec{\gamma} \cdot (1, 1, 1, 1, 1) = 0 \end{cases}$$

i.e.

$\vec{\gamma} \in \mathbf{ker}(\mathbf{A}) \rightarrow$ only 1 dimension left (normalization)

Nitrogen dioxide

The classical Brewer algorithm

$$\begin{pmatrix} \sigma_{O_3}(\lambda_1) & \sigma_{O_3}(\lambda_2) & \sigma_{O_3}(\lambda_3) & \sigma_{O_3}(\lambda_4) & \sigma_{O_3}(\lambda_5) \\ \beta_{Ray}(\lambda_1) & \beta_{Ray}(\lambda_2) & \beta_{Ray}(\lambda_3) & \beta_{Ray}(\lambda_4) & \beta_{Ray}(\lambda_5) \\ \delta_{aer}(\lambda_1) & \delta_{aer}(\lambda_2) & \delta_{aer}(\lambda_3) & \delta_{aer}(\lambda_4) & \delta_{aer}(\lambda_5) \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \gamma_4 \\ \gamma_5 \end{pmatrix} = \mathbf{0}_{K_{4,1}}$$

i.e.


$$\begin{cases} \vec{\gamma} \perp \vec{\sigma}_{O_3} \\ \vec{\gamma} \perp \vec{\beta}_{Ray} \\ \vec{\gamma} \perp \vec{\delta}_{aer} \\ \vec{\gamma} \cdot (1, 1, 1, 1, 1) = 0 \end{cases}$$

i.e.

$\vec{\gamma} \in \mathbf{ker}(\mathbf{A}) \rightarrow$ only 1 dimension left (normalization)

Nitrogen dioxide

Algorithm improvements



Nitrogen dioxide

Algorithm improvements

We need additional degrees of freedom to remove influence of all absorbers

- add 1 slit and search for $\vec{\gamma}$ so that $|\vec{\gamma} \cdot \sigma_{\vec{NO}_2}| = \max$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{O}_4}$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{H}_2O}$
- search for a new operating step of micrometer \rightarrow new set of wavelengths (all past data are lost)

Nitrogen dioxide

Algorithm improvements

We need additional degrees of freedom to remove influence of all absorbers

- add 1 slit and search for $\vec{\gamma}$ so that $|\vec{\gamma} \cdot \sigma_{\vec{NO}_2}| = \max$
- or add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{O}_4}$
- or add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{H}_2O}$
- search for a new operating step of micrometer \rightarrow new set of wavelengths (all past data are lost)

Nitrogen dioxide

Algorithm improvements

We need additional degrees of freedom to remove influence of all absorbers

- add 1 slit and search for $\vec{\gamma}$ so that $|\vec{\gamma} \cdot \sigma_{\vec{NO}_2}| = \max$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{O}_4}$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{H}_2O}$
- search for a new operating step of micrometer \rightarrow new set of wavelengths (all past data are lost)

Nitrogen dioxide

Algorithm improvements

We need additional degrees of freedom to remove influence of all absorbers

- add 1 slit and search for $\vec{\gamma}$ so that $|\vec{\gamma} \cdot \sigma_{\vec{NO}_2}| = \max$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{O}_4}$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{H}_2O}$
- search for a new operating step of micrometer \rightarrow new set of wavelengths (all past data are lost)

Nitrogen dioxide

Algorithm improvements

We need additional degrees of freedom to remove influence of all absorbers

- add 1 slit and search for $\vec{\gamma}$ so that $|\vec{\gamma} \cdot \sigma_{\vec{NO}_2}| = \max$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{O}_4}$
- **or** add 1 slit and search for $\vec{\gamma}$ so that $\vec{\gamma} \perp \sigma_{\vec{H}_2O}$
- search for a new operating step of micrometer \rightarrow new set of wavelengths (all past data are lost)

Nitrogen dioxide

Algorithm improvements

Alternative approach: **jump scans** → increase the number of equations and determine more unknowns:

- NO_2
- O_3
- O_4
- H_2O
- Rayleigh
- aerosol optical depth (AOD)
- Ring effect
- effective NO_2 temperature
- wavelength misalignments

Best if overconstrained system → SVD / least squares solution

Nitrogen dioxide

Algorithm improvements

Alternative approach: **jump scans** → increase the number of equations and determine more unknowns:

- NO_2
- O_3
- O_4
- H_2O
- Rayleigh
- aerosol optical depth (AOD)
- Ring effect
- effective NO_2 temperature
- wavelength misalignments

Best if overconstrained system → SVD / least squares solution

Nitrogen dioxide

Algorithm improvements

Alternative approach: **jump scans** → increase the number of equations and determine more unknowns:

- NO_2
- O_3
- O_4
- H_2O
- Rayleigh
- aerosol optical depth (AOD)
- Ring effect
- effective NO_2 temperature
- wavelength misalignments

Best if overconstrained system → SVD / least squares solution

Nitrogen dioxide

Future developments

- calibration of Brewer #066 (Langley plot)
- determination of air mass factors (AMFs) for direct sun (DS) and zenith sky (ZS) measurements
- → need of photochemical models?
- comparison between the two viewing geometries

- 1 Introduction
- 2 Nitrogen dioxide
- 3 Aerosol properties**
- 4 Conclusions

Aerosol properties

POM02 photometer




Aerosol properties

- supported by Euroskyrad network (NRT processing, quality control, comparison to other sites, etc.)
- several wavelengths (315, 340, 380, 400, 500, 675, 870, 940, 1020, 1627 and 2200 nm);
- two different collimators, but same detector;
- it is possible to retrieve many aerosol properties (which can also be useful for radiative transfer issues)
 - ▶ aerosol optical depth (AOD);
 - ▶ single scattering albedo;
 - ▶ real and imaginary refractive index;
 - ▶ Angstrom exponent;
 - ▶ distribution size
 - ▶ phase function


Aerosol properties

POM02 photometer




Aerosol properties

First results (Brewer)



Aerosol properties

First results



Aerosol properties

First results




Aerosol properties

First results




Aerosol properties

First results



Aerosol properties

First results



- 1 Introduction
- 2 Nitrogen dioxide
- 3 Aerosol properties
- 4 Conclusions**

Conclusions

- photometric techniques cannot replace local measurements;
- i.e. they does not work with cloudy sky;
- however, they represent a complementary approach to air/climate monitoring;
- photometric measurements are representative of a wide area;
- solar monitoring has great importance!